



© J ORTHOP TRAUMA SURG REL RES

16(10) 2021

Research Article

# Colour Doppler vs compression ultrasound as diagnostic modality for acute venous thromboembolism in patient with proximal femur fracture

RAHUL SAKET, ARUN K NAIK, TARINI PRASAD MOHANTY

Department of Orthopaedics, Hi-Tech Medical College, Rourkela, Orissa, India

Address for correspondence:

Rahul Saket, Department Of Orthopaedics, Hi-Tech Medical College, Rourkela, Orissa, India

drrsinha3@gmail.com

## Statistics

Figures	00
Tables	04
References	32

Received: 07.10.2021

Accepted: 01.11.2021

Published: 15.11.2021

## Abstract

**Introduction:** This research is a comparative study between colour Doppler and compression ultrasound, used as diagnostic modalities for diagnosing DVT in patients with proximal femur fracture.

**Material and Methods:** It was a prospective study conducted over a period of one and a half years. Cases under study were patients with isolated proximal femur fracture. All the patients were evaluated by colour Doppler and compression ultrasound. CT Angiography was used whenever required and not as routine modality. Post-operative monitoring of the patients was done for 8 weeks.

**Results:** A total of 70 patients were assessed. The mean age group was 50 years, with 60% being males and 40% females. There were 10 cases of DVT diagnosed radiologically. Out of these 10 cases diagnosed, 9 cases were diagnosed positive by the colour Doppler method and 8 by the compression ultrasound. One case was diagnosed by CT Venography. There was complete resolution of DVT in all the patients who were diagnosed positive for DVT. It was confirmed by colour Doppler at 8 weeks post diagnosis.

**Conclusion:** DVT is a serious complication if untreated. Previous studies showing lesser efficacy of colour Doppler as compared to compression ultrasound does not hold true. Cases diagnosed by colour Doppler are at par with compression ultrasound. So we can use colour Doppler as screening modality with the added benefit of visualization of thrombus resolution

**Keywords:** deep vein thrombosis, venous thromboembolism, pulmonary thrombo-embolism

## INTRODUCTION

Venous Thromboembolism (VTE) is a common preventable cause of morbidity and mortality in trauma patients who survive the first 24 hours. It includes Deep Venous Thrombosis (DVT) in extremities, pelvic veins and their embolization to the Pulmonary Circulation (PE). Occurrence of DVT in trauma patients varies, ranging from 5% to 70% depending on the magnitude of injury and modalities of diagnosis [1-5]. DVT manifests clinically with pain and swelling of the lower limb. Pulmonary embolism manifests with respiratory distress and death. Many times clinical signs may be minimal or absent and pulmonary embolism may be the very first manifestation [1,3]. But in cases of trauma, many symptoms are confusing like swelling/pain in lower limb, respiratory distress, etc., which can be due to trauma itself [4]. Proximal femur fractures result from high velocity trauma in younger age groups and due to osteoporosis in older age groups. Trauma is not only associated with fractures but also with damage to vascular structure around the hip [1,6].

Studies on Indian patients reported variable incidence of DVT (0%-46%) [7-11]. One of the reasons for this variation in results may be the different diagnostic modalities used in these cases. With CT Venography, high incidence was reported [7]. One of the reasons being that CT Venography diagnoses minute thrombi, which are clinically non-significant. But CT Venography cannot be used as routine modality because of high radiation exposure, high cost and the infrastructure required.

In routine we use colour Doppler or compression ultrasound for diagnosing DVT. Colour Doppler is the newer modality with benefits to identify non-occlusive thrombi/resolving thrombi and to visualize the calf veins that are difficult to assess with conventional ultrasound scanners [12-16]. Information on the accuracy of colour Doppler imaging in symptomless postoperative patients is sparse and limited to 5 studies that provide conflicting results, (with reported sensitivities of 38% to 100%) [17-21]. But there are very few comparative studies between colour Doppler and compression ultrasound for diagnosing DVT. These studies were conducted on western patients either undergoing arthroplasty surgery or on non-trauma patients [22]. These studies concluded that compression ultrasound was better in diagnosing DVT as compared to colour Doppler. So we conducted a comparative study between colour Doppler and compression ultrasound in diagnosing DVT in trauma patients having proximal femur fracture only.

## MATERIALS AND METHODS

It was a prospective study conducted for 18 months at Hi-Tech medical college and hospital. The study was conducted on 70 patients, admitted with isolated proximal femur fracture (neck of femur, inter trochanteric and sub trochanteric fractures) (Table 1). Patients included in this study were between 18 to 65 years of age. Patients excluded from this study were those on whom CT Venography was contraindicated due to renal problem or reaction to the dye used for angiography. Patients having other medical diseases like history of cardiac or respiratory failure, undergoing treatment for malignancy, etc. were excluded from the study as they were on medication which would alter their coagulation profile. Patients on hormonal contraceptives were also excluded from the study. We tried to focus on pure orthopedic patients.

In all the patients, trauma protocol of airway and breathing and circulation monitoring was done. Patients were stabilized. After getting informed consents, patients were enrolled for the study. Routine investigations were done like complete blood count, kidney function test, liver function test, cougulogram, etc. Records were maintained regarding duration of hospital stay, fracture classification, type of anesthesia (general or regional) and surgical approach used. All patients were encouraged for bed side physiotherapy as early as possible, post-surgery. No chemoprophylaxis was used for DVT. On diagnosis of DVT, patients were immediately put on treatment with oral Rivaroxaban.

Bilateral ultrasound examinations were performed by experienced physicians and technicians using a 5 MHz or 7.5 MHz ultrasound transducer. All the patients were first evaluated by compression ultrasound followed by colour Doppler assessment performed by different persons. The examination was started by evaluating the external iliac vein and femoral vein followed by popliteal veins. The upper halves of the peroneal veins and the posterior tibial veins were then examined [23,24] (the examination of the calf veins was limited to the upper half because the distal segments of the calf veins are technically difficult to assess). Veins were always identified by their accompanying arteries. For compression ultrasound examination, the only criterion for the diagnosis of DVT was the inability to compress the veins with the ultrasound probe [23-25]. Patients were considered not to have DVT if all of the veins were fully compressible and no residual lumen was seen. Vein compressibility was always assessed in the transverse view because when compression is performed in the longitudinal plane, the transducer may occasionally slide off the vein and so falsely simulate compressibility. Ultra-sonographic findings were recorded as negative, positive, or inadequate for interpretation. Images in the longitudinal axis were used for the evaluation of the colour Doppler images. At selected sites, transverse views also were obtained. Augmentation of flow was achieved by manual compression of the calf to improve visualization of the calf veins. The criteria for an abnormal colour Doppler test were the absence of colour in a vein after augmentation or a focal intraluminal filling defect. The compression ultrasound and Colour Doppler findings were considered inadequate for interpretation if a complete vein or a segment of a vein could not be identified.

Compression ultrasound and colour Doppler study was done on the 1<sup>st</sup> day of admission, on the day of discharge or day 5 of admission. Then in the 2<sup>nd</sup>, 4<sup>th</sup>, 6<sup>th</sup> and 8<sup>th</sup> weeks post-operatively, during out door follow ups. In cases with doubtful lesions or inadequate interpretation of colour Doppler or compression ultrasound, CT venography was done.

**Table 1.** Type of fracture and number of cases

Type of fracture	Number of cases
Inter-trochantric (I/T)	28
Neck of femur	14
Sub trochantric (S/T)	12
I/T with S/T	16
Total	70

**Table 2.** Diagnostic Modality

Diagnostic modality	Location of thrombus		
	Proximal (Total case= 4)	Proximal and Distal (Total case= 3)	Distal (Total case= 3)
Compression ultrasound	3	3	2
Color Doppler	4	3	2
CT Venography	-	-	1

## RESULTS

A total of 70 patients with fracture proximal femur were assessed and evaluated for 18 months. Out of 70 patients, in 3 patients, colour Doppler and compression ultrasound reading was inadequate, so we carried out CT Venography. Out of 3 patients, 1 case was DVT positive, but thrombus was minute, in distal segment and not clinically significant. A total number of cases that were radiologically proved positive for DVT were 10 (14%) (Table 2).

Positive cases diagnosed by compression ultrasound were eight. Nine cases were diagnosed by colour Doppler. 8 cases were same for both compression ultrasound and colour Doppler (Table 1). A new case of DVT, diagnosed by colour Doppler had thrombus just proximal to trifurcation of popliteal vein. We were not able to comment on the sensitivity and specificity as CT Venography was not done on all patients. 1 case was diagnosed DVT positive in the 5<sup>th</sup> week of follow up (Table 3).

Out of the total cases the number of proximal DVT was 4, the number of distal DVT was 3 and 3 cases had both proximal and distal DVT (Table 2). Cases of proximal DVT, diagnosed by compression ultrasound were 3 and by colour Doppler, 4. The number of cases having distal DVT was 3, of which 2 was diagnosed bysonological method and 1 case by CT Venography. 3 cases had both proximal and distal DVT and were diagnosed by both compression ultrasound and colour Doppler (Table 2). Better diagnosis of distal DVT by CT Venography was anticipated as colour Doppler and compression ultrasound are not modalities for distal thrombus. But distal thrombus has lesser probability of pulmonary embolism.

No case of DVT was reported patients in the age group of 18-30 years (Table 4). In the age group of 31-50 years, 4 patients were diagnosed positive. In the age group of 51-65, 6 patients were diagnosed positive. However the difference in age groups was not statistically significant. All patients were operated under regional anesthesia. In most of the patients, we went for osteosynthesis except for the 10 geriatric patients of fracture neck femur, where we went for hemi-arthroplasty.

In the post-operative period there was no case of surgical site infection. Suture removal was done on the 14<sup>th</sup> day, post-operative. Physiotherapy protocol included non-weight bearing to partial and full weight bearing, crutch walking, active and passive hip and knee range of movement, side sitting, etc.

All cases of radiologically proven DVT were treated with Tab Rivaroxaban (15 mg, BD for 3 weeks then 20 mg OD for further 3 weeks). Thrombus resolution was assessed by normal venous flow on Doppler study. In all patients of DVT, complete resolution of thrombi was noted at 8 weeks.

**Table 3.** Number of DVT Cases

Time post injury	Number of DVT cases
Within 1 week	3
1-2 week	2
2-4 week	4
4-6 week	1
Total	10

**Table 4.** Patient data and DVT cases

Age group	Number of patient	DVT cases
18-30	8	0
30-50	28	4
50-65	34	6
Total	70	10

## DISCUSSION

DVT is a common but preventable complication in trauma patients [1,3]. Indians are supposed to have low incidence of DVT. However, the studies on orthopedic patients in India gave variable incidence [7-11].

Till date all studies on Indian patients have used a single diagnostic modality. High incidence was noted when CT Venography was used [7]. Normally we use colour Doppler and compression ultrasound as routine diagnostic modalities. On searching PubMed and google, we found only one comparative study between colour Doppler and compression ultrasound [22]. This study was on western patients, undergoing arthroplasty surgery. This study showed better efficacy of compression ultrasound. Ongoing through few other related articles [26], general inference was that colour Doppler is less efficient for diagnosing DVT. But our study contradicts this notion. In our study we found colour Doppler at par with compression ultrasound.

By studying on Indian patients, we were able to clear several myths. With regard to present aim of study being related to diagnostic modality, colour Doppler is at par with compression ultrasound. Though cases diagnosed by colour Doppler were more, but considering the sample size we were not able to give generalized statement of better efficacy of colour Doppler. Since the population in our country is large, so we generally choose single diagnostic modality. Previously it was compression ultrasound, but now we can shift to colour Doppler for better results. Colour Doppler can even better visualize resolution of thrombus.

Another fact which we found in our study was that DVT can occur up to 4 weeks post operation [5,27,28]. This is an important concern. In all the previous studies, results were based on indoor patients. But the risk of DVT is there even in the follow up period. All previous studies undertook baseline CT Venography scanning of all patients. Since DVT can occur in follow up period so it is not possible ethically as well as practically to expose patients for CT in each follow up.

Another significance of our study was that we conducted our study on trauma patients. Trauma patients require more thoughtful evaluation because pathogenesis of DVT starts immediately after injury. Many other factors are responsible for hypercoagulability [29] in post trauma patients like lack of adequate hydration, prolonged immobilization [1,5], decreased serum levels of Ant thrombin III [29,30] and damage to soft tissue. Surgical procedure adds to further insult. Trauma is itself associated with lower limb swelling giving false positive clinical impression.

Our study had a few weaknesses that must be considered. First of all sample size was small. Secondly, we were not able to document on sensitivity and specificity as we did not carry out baseline CT Venography of all patients. It would have given more authenticity to our study, but we were more focused on screening modality.

Since there are only two studies comparing colour Doppler and compression ultrasound (including ours), so more study in this regard is required. This will help us in formulating screening protocol for DVT. Screening protocol consists of schedule of screening, modality to be used and requirement of CT venography. DVT screening should also be done in follow up period, not only in indoor period, as DVT can occur in post-operative period. DVT monitoring is much debated in the Indian subcontinent as DVT incidence is considered low here and same result is superimposed on trauma patients.

## CONCLUSION

In our study we found that result of colour Doppler is better or at par with compression ultrasound in diagnosing DVT. Colour Doppler has

added benefit of visualization of thrombus resolution. Like compression ultrasonography, colour Doppler ultrasound has limitations for screening DVT in calf muscle and beyond. Further refinements in ultrasound technology can improve the functional outcome. In particular, increased technical sensitivity to faint flow signals such as power Doppler [31] and the application of ultrasound contrast agents show promise.

## References:

1. Geerts W.H., Code K.I., Jay R.M., et al.: A prospective study of venous thromboembolism after major trauma. *N Engl J Med.* 1994;331:1601-1606.
2. Montgomery K.D., Geerts W.H., Potter H.G., et al.: Thromboembolic complications in patients with pelvic trauma. *Clin Orthop Relat Res.* 1996;329:68-87.
3. Napolitano L.M., Garlapati V.S., Heard S.O., et al.: Asymptomatic deep venous thrombosis in the trauma patient: is an aggressive screening protocol justified? *J Trauma.* 1995;39:651-657.
4. Knudson M.M., Collins J.A., Goodman S.B., et al.: Thromboembolism following multiple trauma. *J Trauma.* 1992;32:2-11.
5. Geerts W.H., Bergqvist D., Pineo G.F., et al.: Prevention of venous thromboembolism: American college of chest physicians evidence-based clinical practice guidelines (8<sup>th</sup> edition). *Chest.* 2008;133:381S-453S.
6. Montgomery K.D., Geerts W.H., Potter H.G., et al.: Practical management of venous thromboembolism following pelvic fractures. *Orthop Clin N Am.* 1997;28:397-404.
7. Agarwala S., Bhagwat A.S., Modhe J., et al.: Deep vein thrombosis in Indian patients undergoing major lower limb surgery. *Indian J Surg.* 2003;37:159-62.
8. Sharma H., Maini L., Agrawal N., et al.: Incidence of deep vein thrombosis in patients with fractures around hip joint: a prospective study. *Indian J Orthop.* 2002;36:5.
9. Agarwala S., Bhagwat A.S., Wadhvani R., et al.: Pre and postoperative DVT in Indian patients- Efficacy of LMWH as a prophylaxis agent. *Indian J Orthop.* 2005;39:55e58.
10. Maini P.S., Talwar N., Nijhawan V.K., et al.: Results of cemented bipolar hemiarthroplasty for fracture of the femoral neck-10 year study. *Indian J Orthop.* 2006;40:154e156.
11. Bagaria V., Modi N., Panghate A., et al.: Vaidya S. Incidence and risk factors for development of venous thromboembolism in Indian patients undergoing major orthopaedic surgery: results of a prospective study. *Postgrad Med.* 2006;82:136e139.
12. Foley W.D., Middleton W.D., Lawson T.L., et al.: Colour Doppler ultrasound imaging of lower extremity venous disease. *AJR Am J Roentgenol.* 1989;152:371-376.
13. Persson A.V., Jones C., Zide R., et al.: Use of the triplex scanner in diagnosis of deep venous thrombosis. *Arch Surg.* 1989;124:593-597.
14. Baxter G.M., McKechnie, Duffy P., et al.: Colour Doppler ultrasound in deep venous thrombosis: a comparison with venography. *Clin Radiol.* 1990;42:32-36.
15. Rose S.C., Zwiebel W.J., Nelson B.D., et al.: Symptomatic lower extremity deep venous thrombosis: accuracy, limitations, and role of colour duplex flow imaging in diagnosis. *Radiology.* 1990;175:639-644.
16. Schindler J.M., Kaiser M., Gerber A., et al.: Colour coded duplex sonography in suspended deep vein thrombosis of the leg. *BMJ.* 1991;301:1369-1370.
17. Cronan J.J., Froehlich J.A., Dorfman G.S., et al.: Image-directed Doppler ultrasound: a screening technique for patients at high risk to develop deep vein thrombosis. *J Clin Ultrasound.* 1991;19:133-138.
18. Davidson B.L., Elliott G.C., Lensing A.W.A., et al.: Low accuracy of colour Doppler ultrasound in the detection of proximal leg vein thrombosis in asymptomatic high risk patients. *Ann Intern Med.* 1992;171:735-738.
19. Mattos M.A., Londey G.L., Leutz D.W., et al.: Colour-flow duplex scanning for the surveillance and diagnosis of acute deep venous thrombosis. *J Vasc Surg.* 1992;15:366-376.
20. Nicolaidis A., Kalodiki E.: Duplex scanning in post-operative surgical patients. *Haemostasis.* 1993;(suppl 1):72-79.
21. Magnusson M., Erikson B.L., Kalebo P., et al.: Is colour Doppler ultrasound a sensitive screening method in diagnosing deep vein thrombosis after hip surgery? *Thromb Haemost.* 1996;75:242-245
22. Lensing A.W., Doris C.I., McGrath F.P., et al.: A comparison of compression ultrasound with colour Doppler ultrasound for the diagnosis of symptomless postoperative deep vein thrombosis. *Arch Intern Med.* 1997;157:765-8.
23. Lensing A.W.A., Prandoni P., Brandjes D., et al.: Detection of deep vein thrombosis by real-time B-mode ultrasonography. *N Engl J Med.* 1989;320:342-345.
24. White R.H., McGahan J.P., Dasenbach M.M., et al.: Diagnosis of deep vein thrombosis using duplex ultrasound. *Ann Intern Med.* 1989;111:297-303.
25. Heijboer H., Buller H.R., Lensing A.W.A., et al.: A comparison of real-time compression ultrasonography with impedance plethysmography for the diagnosis of deep vein thrombosis in symptomatic outpatients. *N Engl J Med.* 1993;329:1365-1369.
26. Planes A., Vochelle N., Darman J.Y., et al.: Risk of Deep-vein thrombosis after hospital discharge in patients having undergone total hip replacement. *Lancet.* 1996; 348:224-8.
27. Dahl O.E., Andressen G., Aspelin T., et al.: Prolonged prophylaxis following Hip replacement surgery, *J Thromb Haemostat* 1997;77:26-31.
28. Selby R., Geerts W., Ofose F.A., et al.: Hypercoagulability after trauma: hemostatic changes and relationship to venous thromboembolism. *Thromb Res.* 2009;124:281e287.
29. Miller R.S., Weatherford D.A., Stein D., et al.: Antithrombin III and trauma patients: factors that determine low levels. *J Trauma.* 1994 Sep;37:442e445.
30. Rubin J.M., Bude R.O., Carson P.L., et al.: Power Doppler US: a potentially useful alternative to mean frequency-based colour Doppler US. *Radiology.* 1994;190: 853-856.
31. Balen F.G., Allen C.M., Lees W.R., et al.: Ultrasound contrast agents. *Clin Radiol.* 1994; 49:77-82.

## ACKNOWLEDGEMENTS

I am highly grateful to my team members for their co-operation in my work. My colleague Dr. Arun Naik has helped me a lot in patient screening and in follow up of patients. My junior resident Dr. Tarini helped in patient counselling and in data tabulation. I am also grateful to my institution for allowing me to carry out our work.